

# The ultrastructure and immunohistochemistry of the septum pellucidum in a case of thalamic low grade astrocytoma with review of literature

This study was presented in the Symposium Internationale Quartum Anatomiae Clinicae 2000, Varna, Bulgaria as a poster presentation.

M. Mustafa Aldur (1)  
Mustafa Berker (2)  
H. Hamdi Celik (1)  
Mustafa F. Sargon (1)  
Yesim Ugur (3)  
Attila Dagdeviren (3)

(1) Department of Anatomy, Hacettepe University, Faculty of Medicine, Ankara, Turkey  
(2) Department of Neurosurgery, Hacettepe University Faculty of Medicine, Ankara, Turkey  
(3) Department of Histology & Embryology, Hacettepe University Faculty of Medicine, Ankara, Turkey

## Correspondence Address

M. Mustafa Aldur, MD, PhD  
Hacettepe University Faculty of Medicine  
Department of Anatomy, 06100, Ankara, Turkey  
Phone : 90 (312) 305 24 66  
Fax : 90 (312) 305 24 66  
e-mail : mustafa@aldur.net

## Abstract

This article presents the results of light and electron microscopic (EM) findings of the septum pellucidum (Sp) in a case of thalamic low grade astrocytoma. Furthermore, it presents immunohistochemical staining of the Sp which has not been previously reported in the literature.

The tissue specimen was obtained from a 34 year-old male patient, who had undergone subtotal tumor excision operation by the neurosurgery department. The intact part of the Sp was taken for routine light, transmission (TEM), scanning electron microscopic (SEM) and immunohistochemical examinations.

In light microscopy and TEM examination the myelinated axons were small and medium sized. There was no grey matter in the examined area. In SEM examination, there were holes between the ependymal cells that showed transition zones between the inside of cavity and outside of it. In the immunohistochemical examination of the cryosections of the Sp, the ependymal lining cells were reactive with mAb B-F45. Significance and function of this antigen on ependymal cells remains to be clarified. The connection, relationships and function of the Sp is important in the human body. This detailed case study gives information for further research.

Received 10 September 2002; accepted 27 September 2002

**Key words:** septum pellucidum, ultrastructure, immunohistochemistry, astrocytoma

## Introduction

The septum pellucidum (septum gliosum [1-3], septum lucidum [4], postcommissural septum [5], supracommissural septum [6], Sp) is usually described as a thin (1.5-3 mm [7]), vertical partition composed of two laminae, largely separated by a narrow cavity (ventricule de la cloison [4], fifth ventricle [8-9], cavum septi pellucidi), an extrapial space, and therefore, not communicating with the brain's ventricular system. Seen from aside, the Sp is in the shape of a triangle, its base located anteriorly and apex posteriorly. It is attached above to the lower surface of the trunk of the corpus callosum (Cc), below and behind to the anterior part of the fornix, below and in front to the upper surface of rostrum of the Cc, and directly in front to the genu of the Cc. Each lamina forms part of the medial wall of the anterior horn and central part of the lateral ventricle, being therefore lined by ependymal cells on its ventricular side and pia on its medial aspect [10]. In most cases the two laminae are found to be attached to each other in midline, without a cavum septi pellucidi (CaSp) [11]. Classically each lamina of the Sp is made of neural tissue, a thin outer layer of gray matter, and an inner layer of white matter [12-13]. The laminal gray matter (ganglion du septum lucidum [4], nucleus septofimbrialis) is a dorsocaudal extension of the lateral septal nucleus which is located in precommissural septum (area parolfactoria of Broca, area

precommissuralis, area paracommissuralis [5], septum verum [14], septum gangliosum [1-2]) [5]. Some antigenic and ultrastructural aspects of this structure are still unclear.

In this case study, first, we examined the light microscopic, and electron microscopic (transmission, scanning) appearance of the septum pellucidum. Secondly, we performed immunohistochemical staining which has not been previously reported in the literature.

## Material and Methods

The tissue specimen was obtained from a 34 year-old male patient, who had undergone subtotal tumor excision operation by the neurosurgery department. This patient had a thalamic low grade astrocytoma and during the operation, transcortical – transventricular approach was performed (Figs.1a, b). Additionally, in the operation, a fenestration was performed to the septum pellucidum because of the hydrocephalus. The fenestrated part of the septum pellucidum was taken for routine light, transmission, scanning electron microscopic and immunohistochemical examinations.

The tissue specimen was immediately frozen in liquid nitrogen and stored at -24°C. Cryo sections (5-7µm thick) were taken and placed on gelatin-coated slides and kept in humidity-free containers at room temperature, and the staining was carried out within a few days.




Figure 1a MR image of the patient with thalamic low grade astrocytoma. Arrow indicates the tumoral mass. Asterisk indicates the intact part of the septum pellucidum.




Figure 1b MR image of the patient with thalamic low grade astrocytoma. Arrow indicates the tumoral mass. Asterisk indicates the intact part of the septum pellucidum.

Primary monoclonal antibodies (mAb) used in this study, (originator Vermont-Desroches) were supplied from the Seventh HLDA Workshop Endothelial Panel, Harrogate 2000, England.

The indirect immunoperoxidase procedure used in the study has been described in detail by Dijkstra et al. [15]. Sections were fixed in acetone for 10 minutes and air-dried for at least 30 minutes. Sections were then incubated for 60 minutes with mAbs. After washing in 0.01 M phosphate buffered saline (PBS) pH 7.4, the sections were covered with rabbit anti-mouse Ig-G peroxidase conjugate (Sigma Cat No: A-9044, St. Louis) in PBS (dilution 1:200), containing 22% bovine serum albumin (Sigma Cat No: A-7034, St. Louis) (dilution 2.5:100) and normal human serum (dilution 1:100). After washing in PBS the slides were stained for peroxidase activity with 3,3'-diaminobenzidine-tetrahydrochloride (Sigma Cat No:

D-5637, St. Louis) in 0.05 M Tris-HCL buffer pH 7.6 containing H<sub>2</sub>O<sub>2</sub>. Counterstaining was done with haematoxylin. Dual control staining was performed by omitting the first (primary antibody) step and using a mouse IgG. Stained sections were examined and photographed using Olympus BH2 microscope.

For light and transmission electron microscopy, the specimens were fixed in 2.5% glutaraldehyde for 24 hours, washed in phosphate buffer (pH: 7.4), post-fixed in 1% osmium tetroxide in phosphate buffer (pH: 7.4) and dehydrated in increasing concentrations of ethanol. Then, the tissues were washed with propylene oxide and embedded in epoxy-resin embedding media. Semi-thin sections about 2 μm in thickness and ultrathin sections about 60nm in thickness were cut with a glass knife on a LKB-Nova (Sweden) ultramicrotome. Semi-thin sections were stained with methylene blue and examined by a Nikon




Figure 2 Myelinated axons (MA) in the light microscopic examination of Sp (X40).




Figure 3 Myelinated axons (MA), and glial cells (GC) in TEM examination of Sp. (X5000).




Figure 4a Regular myelinated axons (MA) in the SEM examination of Sp (X5000).




Figure 4b Glial cells (GC) between the myelinated axons in the SEM examination of Sp (X5000).




Figure 4c Endepymal cells (EC) in the SEM examination of Sp (X200).




Figure 4d The hole (H) between the endepymal cells (X400).

Optiphot (Japan) light microscope. Ultrathin sections were collected on copper grids, stained with uranyl acetate and lead citrate and examined with a Jeol JEM 1200 EX (Japan) transmission electron microscope. For scanning electron microscopy, fixed samples were dehydrated in increasing concentrations of acetone and critical point dried, mounted on metal stubs with conductive silver paint and then sputtered with a 10 nm thick layer of gold in a BIO-RAD sputter apparatus (England). The tissue samples were examined with a Jeol scanning electron microscope (SEM ASID-10, Japan) at an acceleration voltage of 80 kV.

## Results

In the light and transmission electron microscopic examination, myelinated axons, glial cells and blood vessels were observed. These myelinated axons were small and medium sized (Figs. 2, 3). The parts of the Sp which were examined did not contain any grey matter and for this reason no neuronal structures were found in these examined areas.


In the SEM examination, a large amount of myelinated axons were observed (Fig. 4a). The myelin composition in those axons were regular. Additionally,

in some areas, glial cells were found around the myelinated axons (Fig. 4b). In addition to these structures endepymal cells were presented in the scanning electron microscopic examination (Fig. 4c). The most important finding observed in SEM was the holes found in between the endepymal cells (Fig. 4d). These holes were 10-60µm in diameter. The presence of these holes showed the transition zones between the inside of cavity and outside of it. In the immunohistochemical examination of the cryosections of septum pellucidum, by indirect immunoperoxidase technique, the endepymal lining cells were reactive with mAb B-F45 (Fig. 5).

## Discussion

According to some authors, the Sp is a part of the limbic system [16-21]. The direct stimulation of the Sp causes the contractions of the urinary bladder [22]. In case of immunologic blockage of the Sp with antiseptal antibodies schizophrenic manifestations can occur [23]. Additionally, there are a lot of information about its function [19, 24-28], connections [5, 20, 24], and relationships with psychiatric disorders [16-17, 28-35]. The antigenic structure of septum

pellucidum obtained by using immunohistochemical techniques is very important in psychiatric disorders. Our study shows the importance of full-panel staining in larger series. The antigen recognized by B-F45 mAb is yet undefined. It is reactive with various endothelial cell types (artery, vein, high endothelial venule and umbilical vessel endothelium) and other cell groups including cells of perineurium in peripheral nerves, synovial cells and some fibroblasts in different organs examined so far (our unpublished observation). Significance and function of this antigen on ependymal cells remains to be clarified. No reactivity was observed in the immunohistochemical stainings using antibodies CD 227 (BC2 and BC3), CD 146 (MUC18), BD 46 and VJ1/5. However, the staining of ependyma of septum pellucidum by B-F45 was an interesting finding.



The developmental abnormalities of the Sp are CaSp, cavum vergae (CaV), agenesis of the Sp, and a unique case of foramen formation on the Sp [16-17, 33]. It is reported that abnormalities of the Sp, especially CaV are more frequent in women [33]. On the other hand, some authors state that CaSp is more common in men [32]. Although a Sp syndrome caused by its abnormalities is described [20, 33], some authors state that an isolated absence of the Sp will not cause intellectual, neurologic, and behavioral defects [36-37]. The abnormalities of the Sp can be diagnosed prenatally using ultrasonography [38]. The comparison of antigenic structure of septum pellucidum in normal and abnormal cases is very important in the definition of molecular mechanisms of these abnormalities. This case study gives information to further research.

Figure 5 The ependymal lining cells were reactive with mAb B-F45 in the immunohistochemical examination of the serial cryosections of Sp. Arrows indicate the reactivity (X200).

## References

- [1] Andy OJ, Stephan H. The septum in the human brain. *J. Comp. Neurol.* 1968 (133) 383-410.
- [2] Kuhlenbeck H. Some comments on the development of the human corpus callosum and septum pellucidum. *Kaibogaku-Zasshi.* 1969 (44) 245-256.
- [3] Rakic P, Yakovlev PI. Development of the corpus callosum and cavum septi in man. *J. Comp. Neur.* 1968 (132) 45-72.
- [4] Rouvière H, Cordier G, Delmas A. *Anatomie humaine descriptive et topographique, Neuvième Édition Révisée, Tome III, Paris, Masson et Cie, 1962; pp 592-593.*
- [5] Crosby EC, Humphrey T, Lauer EW. *Correlative anatomy of the nervous system.* 1th edn. The MacMillan Company, New York, 1962; pp 350-572.
- [6] Williams PL, Bannister LH, et al. *Gray's anatomy.* 38th. edn. Churchill Livingstone, Edinburgh, 1995; pp 1528-1115-1141.
- [7] Pendergrass EP, Hodes PJ. Dilatation of the cavum septi pellucidi and cavum vergae. *Ann. Surg.* 1935 (101) 269-295.
- [8] Dandy WE. Congenital cerebral cysts of the cavum septi pellucidi (fifth ventricle) and cavum vergae (sixth ventricle). *Arch. Neurol. Psych.* 1931 (25) 44-66.
- [9] Ranson SW, Clark SL. *The anatomy of the nervous system, its development and function.* 10th edn. WB Saunders Company, Philadelphia, 1964; p 337.
- [10] Williams PL, Warwick R, Dyson M, Bannister LH. *Gray's anatomy.* 37th ed, Churchill Livingstone, Edinburgh, 1989, p 1062.
- [11] Carpenter MB. *Neuroanatomy.* 4th ed, Williams and Wilkins, Baltimore, 1991; p 41.
- [12] Davies DV, Coupland RE. *Gray's anatomy.* 34th edn. Longmans, London, 1972; p 1097.
- [13] Truex RC. *Human neuroanatomy.* 6th edn. Williams and Wilkins, Baltimore, 1969; p 457.
- [14] Clemente CD. *Anatomy of the human body by Henry Gray.* 30th American edn. Lea & Febiger, Philadelphia, 1985; pp 1059-1060.
- [15] Dijkstra CD, Döpp EA, Joling P, Kraal G. The heterogeneity of nuclear phagocytes in lymphoid organs: distinct macrophage subpopulations in the rat recognized by monoclonal antibodies ED1, ED2 and ED3. *Immunology.* 1985 (54) 589-599.
- [16] Aldur MM, Celik HH, Sargon MF, Dagdeviren A, et al. Unreported anatomical variation of septum pellucidum. *Clin. Anat.* 1997 (10) 245-249.
- [17] Aldur MM, Gurcan F, Basar R, Aksit MD. Frequency of septum pellucidum anomalies in non-psychotic population: a magnetic resonance imaging study. *Surg. Radiol. Anat.* 1999 (21) 119-123.
- [18] Lancon JA, Haines DE, Lewis AI, Parent AD. Endoscopic treatment of symptomatic septum pellucidum cysts: with some preliminary observations on the ultrastructure of the cyst wall: two technical case reports. *Neurosurgery.* 1999 (45) 1251-1257.
- [19] Noback CR, Demarest JR. *The human nervous system.* 2nd edn. McGraw-Hill Book Company, New York, 1979; p 431.
- [20] Sarwar M. The septum pellucidum: normal and abnormal. *Am. J. Neuroradiol.* 1989 (10) 989-1005.
- [21] Tien RD, Felsberg GJ, Krishnan R, Heinz ER. MR imaging of diseases of the limbic system. *Am. J. Roentgenol.* 1994 (163) 657-665.

- [22] Kabat H, Magoun HW, Ranson SW. Reaction of the bladder to stimulation of points in the forebrain and midbrain. *J. Comp. Neurol.* 1935 (63) 211-239.
- [23] Heath RG, McCarron KL, O'Neil CE. Antiseptal brain antibody in IgG of schizophrenic patients. *Biol. Psychiatry.* 1989 (25) 681-682.
- [24] Curtis BA, Jacobson S, Marcus EM. *An introduction to the neurosciences.* 1. edn. W.B. Saunders Company, Philadelphia, 1972; pp 425-436.
- [25] Sarnat HB, Netsky MG. *Evolution of the nervous system.* Oxford University Press, New York, 1974; p 247-249.
- [26] Smythies JR. *The neurological foundation of psychiatry.* Mosby, St.Louis, 1977; pp 2877-2890.
- [27] Vinken PJ, Bruyn GW. *Handbook of clinical neurology.* Vol 30. Elsevier/North-Holland Biomedical, Amsterdam, 1977; p 299-336.
- [28] Zeman W, King FA. Tumors of the septum pellucidum and adjacent structures with abnormal affective behavior: an anterior midline structure syndrome. *J. Nerv. Ment. Dis.* 1958 (127) 490-502.
- [29] Degreef G, Bogerts B, Falkai P, Greve B, et al. Increased prevalence of the cavum septum pellucidum in magnetic resonance scans and post-mortem brains of schizophrenic patients. *Psychiatry Res.* 1992 (45) 1-13.
- [30] Degreef G, Lantos G, Bogerts B, Ashtari M, et al. Abnormalities of the septum pellucidum on MR scans in first-episode schizophrenic patients. *Am. J. Neuroradiol.* 1992 (13) 835-840.
- [31] DeLisi LE, Hoff AL, Kushner M, Degreef G. Increased prevalence of cavum septum pellucidum in schizophrenia. *Psychiatry Res.* 1993 (50) 193-199.
- [32] Jurjus GJ, Nasrallah HA, Olson SC, Schwarzkopf SB. Cavum septum pellucidum in schizophrenia, affective disorder and healthy controls: a magnetic resonance imaging study. *Psychol. Med.* 1993 (23) 319-322.
- [33] Scott TF, Price TR, George MS, Brillman J, et al. Midline cerebral malformations and schizophrenia. *J. Neuropsychiatry Cli. Neurosci.* 1993 (5) 287-293.
- [34] Shaw CM, Alvord EC. Cava septi pellucidi et vergae: their normal and pathological states. *Brain.* 1969 (92) 213-223.
- [35] Wolf SS, Hyde TM, Weinberger DR. Malformations of the septum pellucidum: two distinctive cases in association with schizophrenia. *J. Psychiatry Neurosci.* 1994 (19) 140-144.
- [36] Brodsky MC, Glasier CM. Optic nerve hypoplasia. Clinical significance of associated central nervous system abnormalities on magnetic resonance imaging. *Arch. Ophthalmol.* 1993 (111) 66-74.
- [37] Williams J, Brodsky MC, Griebel M, Glasier CM, et al. Septo-optic dysplasia: the clinical insignificance of an absent septum pellucidum. *Dev. Med. Child. Neurol.* 1993 (35) 490-501.
- [38] Bronshtein M, Weiner Z. Prenatal diagnosis of dilated cava septi pellucidi et vergae: associated anomalies, differential diagnosis, and pregnancy outcome. *Obstet. Gynecol.* 1992 (80) 838-842.

— N —